

Determination of antibodies against Helicobacter pylori

Antibodies to *Helicobacter pylori* in serum can be detected by a whole range of immunological and serological techniques such as immunoblotting, immunofluorescence , hemagglutination , complement fixation, latex tests, etc. However, **the most widespread method** is undoubtedly **ELISA** - a simple, fast, cheap and reliable technique. However, the specificity and sensitivity of the method is highly dependent on the antigen used - from whole cells, via ultrasonic sonication, glycine extract to purified proteins. In 1989, the isolation of a high molecular weight surface protein (designated HM-CAP, high molecular weight cell-associated protein) was described, with a specificity and sensitivity of 95%.

Serological detection of antibodies to *Helicobacter pylori* is of clinical importance mainly for long-term follow-up after treatment and for monitoring the success of *Helicobacter pylori* eradication . A decrease in IgG after 6 months of treatment to values < 50% shows a specificity of 95% and a sensitivity of 97%. Indications include screening of high-risk patients, e.g. kidney transplant patients , when helicobacter infection increases the risk of peptic ulcer and bleeding.

Antibodies to *Helicobacter pylori* can also be detected in **saliva or urine using immunological techniques**. There are also a number of so-called rapid tests , where antibodies to Helicobacter pylori are detected from whole blood, after capillary sampling within a few minutes using the immunoaffinity chromatography method. The specificity and sensitivity of these tests is relatively low - 70 to 85%.

The serological diagnosis of antibodies to *Helicobacter pylori* also includes the detection of cagA, vacA and iceA antigens, the presence of which specifies strains of *Helicobacter pylori* with higher pathogenicity. Classic ELISA tests on microtitre plates or PCR techniques are used to detect these antigens. Detection of *Helicobacter pylori* and its strains by PCR methods is in the phase of clinical testing, it is not yet used in routine diagnostics. The diagnostic value of determining antibodies to *Helicobacter pylori* is still a subject of research, it is not suitable for screening programs.

SENZITIVITA A SPECIFICITA TESTU NA Hp		
TEST	SENSITIVITA	SPECIFICITA
HISTOLOGIE = 3 BIOPSIE	95	99
RYCHLÝ UREAŽOVÝ CLO TEST	99	99
KULTIVACE	80	99
SEROLOGIE PROTILÁTEK	85	79
PROTILÁTKY VE SLINÁCH	81	73
PROTILÁTKY V MOČI	89	91
RAPID MONOTESTY	71	99
DECHOVÝ TEST = UBT	95	95
DETEKCE ANTIGENU VE STOLICI	99	99

Bures J., Congress-Hadec Křídlové, 1997; Lanza F., Eur.J.Gastro.Hepato.2009
Ohkura R., Scand.J.Gastro. 2009; Vavra D. VAKO N. GUT 2001

Sensitivity and specificity of tests for *H. pylori*

Links

Related Articles

- diagnosis of helicobacter infection
- Breath tests with carbon-13
- Detection of Helicobacter pylori antigen in stool

Source

- KOCNA, Petr. *GastroLab* [online]. [cit. 2011-03-04]. <<http://www1.lf1.cuni.cz/~kocna/glab/glency1.htm>>.

References

- PENG, NJ. Comparison of noninvasive diagnostic tests for Helicobacter pylori infection. *Medical Principles and Practice (International Journal of the Kuwait University Health Sciences Centre)* [online]. 2009, vol. 18, no. 1, p. 57-61, Available from <https://www.ncbi.nlm.nih.gov/pubmed/19060493?ordinalpos=19&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_DefaultReportPanel.Pubmed_RVDocSum>. ISSN 1011-7571, eISSN 1423-0151. PMID: 19060493 (<http://www.ncbi.nlm.nih.gov/pubmed/19060493>).
- JANULAITYTE-GÜNTHER, D. Combined serum IgG response to Helicobacter pylori VacA and CagA predicts gastric cancer. *FEMS immunology and medical microbiology* [online]. 2007, vol. 50, no. 2, p. 220-225, Available from <https://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=17567283&ordinalpos=31&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum>. ISSN 0928-8244, eISSN 1574-695X. PMID: 17567283 (<http://www.ncbi.nlm.nih.gov/pubmed/17567283>).
- ZAMBON, CF. Non-invasive diagnosis of Helicobacter pylori infection: simplified 13C-urea breath test, stool antigen testing, or DNA PCR in human feces in a clinical laboratory setting?. *Clinical biochemistry* [online]. 2004, vol. 37, no. 4, p. 261-267, Available from <<https://www.ncbi.nlm.nih.gov/pubmed/15003727?dopt=Abstract>>. ISSN 0009-9120, eISSN 1873-2933. PMID: 15003727 (<http://www.ncbi.nlm.nih.gov/pubmed/15003727>).
- CHEN, TS. Immunoglobulin G antibody against Helicobacter pylori: clinical implications of levels found in serum. *Clinical and diagnostic laboratory immunology* [online]. 2002, vol. 9, no. 5, p. 1044-1048, Available from <<https://www.ncbi.nlm.nih.gov/pubmed/12204957?dopt=Abstract>>. ISSN 1071-412X, eISSN 1098-6588. PMID: 12204957 (<http://www.ncbi.nlm.nih.gov/pubmed/12204957>).
- KINDERMANN, A. Evaluation of a rapid whole blood test to detect Helicobacter pylori infection in children. *Scandinavian journal of gastroenterology* [online]. 2001, vol. 36, no. 6, p. 572-576, Available from <<https://www.ncbi.nlm.nih.gov/pubmed/11424314?dopt=Abstract>>. ISSN 0036-5521, eISSN 1502-7708. PMID: 11424314 (<http://www.ncbi.nlm.nih.gov/pubmed/11424314>).
- ARENTS, NL. The importance of vacA, cagA, and iceA genotypes of Helicobacter pylori infection in peptic ulcer disease and gastroesophageal reflux disease. *The American journal of gastroenterology* [online]. 2002, vol. 97, no. 4, p. 1065, Available from <<https://www.ncbi.nlm.nih.gov/pubmed/11569682?dopt=Abstract>>. ISSN 0002-9270, eISSN 1572-0241. PMID: 11569682 (<http://www.ncbi.nlm.nih.gov/pubmed/11569682>).